

THE JUNE 2008 FLARE OF MARKARIAN 421 FROM OPTICAL TO TEV ENERGIES

I. DONNARUMMA^{1,*}, V. VITTORINI^{1,2}, S. VERCELLONE³, E. DEL MONTE¹, M. FEROCI¹, F. D'AMMANDO^{1,2}, L. PACCIANI¹, A. W. CHEN^{3,4}, M. TAVANI^{1,2}, A. BULGARELLI⁵, A. GIULIANI³, F. LONGO⁶, G. PUCCELLA¹, A. ARGAN¹, G. BARBIELLINI⁶, F. BOFFELLI⁷, P. CARAVEO³, P. W. CATTANEO⁷, V. COCCO¹, E. COSTA¹, G. DE PARIS¹, G. DI COCCO⁵, Y. EVANGELISTA¹, M. FIORINI³, T. FROYSLAND^{2,4}, M. FRUTTI¹, F. FUSCHINO⁵, M. GALLI⁸, F. GIANOTTI⁵, C. LABANTI⁵, I. LAPSHOV¹, F. LAZZAROTTO¹, P. LIPARI⁹, M. MARISALDI⁵, M. MASTROPIETRO¹⁰, S. MEREGHETTI³, E. MORELLI⁵, A. MORSELLI¹¹, A. PELLIZZONI³, F. PEROTTI³, P. PICOZZA¹¹, G. PORROVECCHIO¹, M. PREST¹², M. RAPISARDA¹³, A. RAPPOLDI⁷, A. RUBINI¹, P. SOFFITTA¹, M. TRIFOGLIO⁵, A. TROIS¹, E. VALLAZZA⁶, A. ZAMBRA³, D. ZANELLO⁹, C. PITTORI¹⁴, P. SANTOLAMAZZA¹⁴, F. VERRECCHIA¹⁴, P. GIOMMI¹⁴, S. COLAFRANCESCO¹⁴, L. SALOTTI¹⁵, (THE AGILE TEAM), M. VILLATA^{16,**}, C. M. RAITERI^{16,**}, W. P. CHEN¹⁷, N. V. EFIMOVA¹⁸, B. JORDAN¹⁹, T. S. KONSTANTINOVA¹⁸, E. KOPELOVA¹⁷, O. M. KURTANIDZE^{20,21,22}, V. M. LARIONOV^{18,23}, J. A. ROS²⁴, A. C. SADUN²⁵, (THE GASP-WEBT TEAM), H. ANDERHUB²⁶, L. A. ANTONELLI²⁷, P. ANTORANZ²⁸, M. BACKES²⁹, C. BAIXERAS³⁰, S. BALESTRA²⁸, J. A. BARRIO²⁸, H. BARTKO³¹, D. BASTIERI³², J. BECERRA GONZÁLEZ³³, J. K. BECKER²⁹, W. BEDNAREK³⁴, K. BERGER³⁴, E. BERNARDINI³⁵, A. BILAND²⁶, R. K. BOCK^{31,32}, G. BONNOLI³⁶, P. BORDAS³⁷, D. BORLA TRIDON³¹, V. BOSCH-RAMON³⁷, T. BRETZ³⁸, I. BRITVITCH²⁶, M. CAMARA²⁸, E. CARMONA³¹, A. CHILINGARIAN⁴⁰, S. COMMICHAU²⁶, J. L. CONTRERAS²⁸, J. CORTINA³⁹, M. T. COSTADO^{33,41}, S. COVINO²⁷, V. CURTEF²⁹, F. DAZZI³², A. DE ANGELIS⁴², E. DE CEA DEL POZO⁴³, R. DE LOS REYES²⁸, B. DE LOTTO⁴², M. DE MARIA⁴², F. DE SABATA⁴², C. DELGADO MENDEZ³³, A. DOMINGUEZ⁴⁴, D. DORNER²⁶, M. DORO³², D. ELSAESSER³⁸, M. ERRANDO³⁹, D. FERENC⁴⁵, E. FERNÁNDEZ³⁹, R. FIRPO³⁹, M. V. FONSECA²⁸, L. FONT³⁰, N. GALANTE³¹, R. J. GARCÍA LÓPEZ^{33,41}, M. GARCZARCZYK³⁹, M. GAUC³³, F. GOEBEL^{31,A}, D. HADASCH³⁰, M. HAYASHIDA³¹, A. HERRERO^{33,41}, D. HÖHNE-MÖNCH³⁸, J. HOSE³¹, C. C. HSU^{31,***}, S. HUBER³⁸, T. JOGLER³¹, D. KRANICH²⁶, A. LA BARBERA²⁷, A. LAILLE⁴⁵, E. LEONARDO³⁶, E. LINDFORS⁴⁶, S. LOMBARDI³², M. LÓPEZ³², E. LORENZ^{26,31}, P. MAJUMDAR³⁵, G. MANEVA⁴⁷, N. MANKUZHIYIL⁴², K. MANNHEIM³⁸, L. MARASCHI²⁷, M. MARIOTTI³², M. MARTÍNEZ³⁹, D. MAZIN³⁹, M. MEUCCI³⁶, M. MEYER³⁸, J. M. MIRANDA²⁸, R. MIRZOYAN³¹, J. MOLDÓN³⁷, M. MOLES⁴⁴, A. MORALEJO³⁹, D. NIETO²⁸, K. NILSSON⁴⁶, J. NINKOVIC³¹, I. OYA²⁸, R. PAOLETTI³⁶, J. M. PAREDES³⁷, M. PASANEN⁴⁶, D. PASCOLI³², F. PAUSS²⁶, R. G. PEGNA³⁶, M. A. PEREZ-TORRES⁴⁴, M. PERSIC^{42,48}, L. PERUZZO³², F. PRADA⁴⁴, E. PRANDINI³², N. PUCHADES³⁹, A. RAYMERS⁴⁰, W. RHODE²⁹, M. RIBÓ³⁷, J. RICO^{49,39}, M. RISSI²⁶, A. ROBERT³⁰, S. RÜGAMER³⁸, A. SAGGION³², T. Y. SAITO³¹, M. SALVATI²⁷, M. SANCHEZ-CONDE⁴⁴, P. SARTORI³², K. SATALECKA^{35,***}, V. SCALZOTTO³², V. SCAPIN⁴², T. SCHWEIZER³¹, M. SHAYDUK³¹, K. SHINOZAKI³¹, S. N. SHORE⁵⁰, N. SIDRO³⁹, A. SIERPOWSKA-BARTOSIK⁴³, A. SILLANPÄÄ⁴⁶, J. SITAREK^{31,34}, D. SOB CZYNSKA³⁴, F. SPANIER³⁸, A. STAMERRA³⁶, L. S. STARK²⁶, L. TAKALO⁴⁶, F. TAVECCHIO²⁷, P. TEMNIKOV⁴⁷, D. TESCARO³⁹, M. TESHIMA³¹, M. TLUCZYKONT³⁵, D. F. TORRES^{49,43}, N. TURINI³⁶, H. VANKOV⁴⁷, A. VENTURINI³², V. VITALE⁴², R. M. WAGNER^{31,***}, W. WITTEK³¹, V. ZABALZA³⁷, F. ZANDANEL⁴⁴, R. ZANIN³⁹, J. ZAPATERO³⁰, (THE MAGIC COLLABORATION), V. ACCIARI^{51,52}, E. ALIU⁵³, T. ARLEN⁵⁴, M. BEILICKE⁵⁵, W. BENBOW⁵¹, S. M. BRADBURY⁵⁶, J. H. BUCKLEY⁵⁵, V. BUGAEV⁵⁵, Y. BUTT⁵⁷, K. BYRUM⁵⁸, A. CANNON⁵⁹, A. CESARINI⁶⁰, Y. C. CHOW⁵⁴, L. CIUPIK⁶¹, P. COGAN⁶², P. COLIN⁶³, W. CUI⁶⁴, M. K. DANIEL⁵⁶, R. DICKERBERG⁵⁵, C. DUKE⁶⁵, T. ERGIN⁵⁷, S. J. FEGAN⁵⁴, J. P. FINLEY⁶⁴, G. FINNEGAN⁶³, P. FORTIN⁶⁶, A. FURNISS⁶⁷, D. GALLI⁶⁴, G. H. GILLANDERS⁶⁰, R. GUENETTE⁶², G. GYUK⁶¹, J. GRUBE^{56,59,***}, D. HANNA⁶², J. HOLDER⁵³, D. HORAN^{58,76}, C. M. HUI⁶³, T. BRIAN HUMENSKY⁶⁸, A. IMRAN⁶⁹, P. KAARET⁷⁰, N. KARLSSON⁶¹, M. KERTZMAN⁷¹, D. KIEDA⁶³, J. KILDEA⁵¹, A. KONOPELKO⁷², H. KRAWCZYNSKI⁵⁵, F. KRENNRICH⁶⁹, M. J. LANG⁶⁰, S. LEBOHEC⁶³, G. MAIER⁶², A. MCCANN⁶², M. MCCUTCHEON⁶², A. MILOVANOVIC⁵⁶, P. MORIARTY⁵², T. NAGAI⁶⁹, R. A. ONG⁵⁴, A. N. OTTE⁶⁷, D. PANDEL⁷⁰, J. S. PERKINS⁵³, A. PICHEL⁷³, M. POHL⁶⁹, K. RAGAN⁶⁴, L. C. REYES⁷⁴, P. T. REYNOLDS⁷⁵, E. ROACHE⁵¹, H. J. ROSE⁵⁶, M. SCHROEDTER⁶⁹, G. H. SEMBROSKI⁶⁴, A. W. SMITH⁵⁸, D. STEELE⁶¹, S. P. SWORDY⁶⁸, M. THEILING⁵¹, J. A. TONER⁶⁰, L. VALCARCEL⁶², A. VARLOTTA⁶⁴, S. P. WAKELY⁶⁸, J. E. WARD⁵⁹, T. C. WEEKES⁵¹, A. WEINSTEIN⁵⁴, D. A. WILLIAMS⁶⁷, S. WISSEL⁶⁸, M. WOOD⁵⁴, B. ZITZER⁶⁴, AND (THE VERITAS COLLABORATION)

Draft version December 8, 2008

ABSTRACT

We present optical, X-ray, high energy ($\lesssim 30$ GeV) and very high energy ($\gtrsim 100$ GeV; VHE) observations of the high-frequency peaked blazar Mrk 421 taken between 2008 May 24 and June 23. A high energy γ -ray signal was detected by AGILE with $\sqrt{TS} = 4.5$ on June 9–15, with $F(E > 100 \text{ MeV}) = 42_{-12}^{+14} \times 10^{-8}$ photons $\text{cm}^{-2} \text{ s}^{-1}$. This flaring state is brighter than the average flux observed by EGRET by a factor of ~ 3 , but still consistent with the highest EGRET flux. In hard X-rays (20–60 keV) SuperAGILE resolved a 5-day flare (June 9–15) peaking at ~ 55 mCrab. SuperAGILE, RXTE/ASM and Swift/BAT data show a correlated flaring structure between soft and hard X-rays. Hints of the same flaring behavior are also detected in the simultaneous optical data provided by the GASP-WEBT. A Swift/XRT observation near the flaring maximum revealed the highest 2–10 keV flux ever observed from this source, of 2.6×10^{-9} erg $\text{cm}^{-2} \text{ s}^{-1}$ (i.e. > 100 mCrab). A peak synchrotron energy of ~ 3 keV was derived, higher than typical values of ~ 0.5 –1 keV. VHE observations with MAGIC and VERITAS on June 6–8 show the flux peaking in a bright state, well correlated with the X-rays. This extraordinary set of simultaneous data, covering a twelve-decade spectral range, allowed for a deep analysis of the spectral energy distribution as well as of correlated light curves. The γ -ray flare can be interpreted within the framework of the synchrotron self-Compton model in terms of a rapid acceleration of leptons in the jet.

Subject headings: BL Lacertae objects: individual (Mrk 421); radiation mechanisms: non-thermal;
X-rays: galaxies; gamma-rays: observations; galaxies: jets

1. INTRODUCTION

Mrk 421 is a nearby blazar ($z = 0.031$) and one of the brightest BL Lac objects given its distance of 134.1 Mpc ($H_0 = 71 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_m = 0.27$, $\Omega_\lambda = 0.73$). It was

observed in γ -rays by EGRET (Lin et al. 1992) and it was the first extragalactic object detected at $E > 500$ GeV (Punch et al. 1992). It belongs to the class of High-energy peaked BL Lac objects (HBLs) (Padovani & Giommi 1995), i. e. radio-loud active galactic nuclei with high radio and optical polarization. Its spectral energy distribution (SED) is double-humped with a first peak usually in the soft to medium X-ray range, and a second one at GeV-TeV energies (Sambruna et al. 1996;

- ¹ INAF/IASF-Roma, I-00133 Roma, Italy
² Dip. di Fisica, Univ. Tor Vergata, I-00133 Roma, Italy
³ INAF/IASF-Milano, I-20133 Milano, Italy
⁴ CIFS-Torino, I-10133 Torino, Italy
⁵ INAF/IASF-Bologna, I-40129 Bologna, Italy
⁶ Dip. Fisica and INFN Trieste, I-34127 Trieste, Italy
⁷ INFN-Pavia, I-27100 Pavia, Italy
⁸ ENEA-Bologna, I-40129 Bologna, Italy
⁹ INFN-Roma La Sapienza, I-00185 Roma, Italy
¹⁰ CNR-IMIP, Roma, Italy
¹¹ INFN Roma Tor Vergata, I-00133 Roma, Italy
¹² Dip. di Fisica, Univ. Dell'Insubria, I-22100 Como, Italy
¹³ ENEA Frascati, I-00044 Frascati (Roma), Italy
¹⁴ ASI Science Data Center, I-00044 Frascati(Roma), Italy
¹⁵ Agenzia Spaziale Italiana, I-00198 Roma, Italy
¹⁶ INAF, Osservatorio Astronomico di Torino, Italy
¹⁷ Institute of Astronomy, National Central University, Taiwan
¹⁸ Astronomical Institute, St. Petersburg State University, Russia
¹⁹ School of Cosmic Physics, Dublin Institute For Advanced Studies, Ireland
²⁰ Abastumani Astrophysical Observatory, Georgia
²¹ Astrophysikalisches Institut Potsdam, Germany
²² Landessternwarte Heidelberg-Königstuhl, Germany
²³ Pulkovo Observatory, Russia
²⁴ Agrupació Astronòmica de Sabadell, Spain
²⁵ Department of Physics, University of Colorado at Denver, CO, USA
²⁶ ETH Zurich, CH-8093 Switzerland
²⁷ INAF National Institute for Astrophysics, I-00136 Rome, Italy
²⁸ Universidad Complutense, E-28040 Madrid, Spain
²⁹ Technische Universität Dortmund, D-44221 Dortmund, Germany
³⁰ Universitat Autònoma de Barcelona, E-08193 Bellaterra, Spain
³¹ Max-Planck-Institut für Physik, D-80805 München, Germany
³² Università di Padova and INFN, I-35131 Padova, Italy
³³ Inst. de Astrofísica de Canarias, E-38200 La Laguna, Tenerife, Spain
³⁴ University of Łódź, PL-90236 Lodz, Poland
³⁵ Deutsches Elektronen-Synchrotron (DESY), D-15738 Zeuthen, Germany
³⁶ Università di Siena, and INFN Pisa, I-53100 Siena, Italy
³⁷ Universitat de Barcelona (ICC/IEEC), E-08028 Barcelona, Spain
³⁸ Universität Würzburg, D-97074 Würzburg, Germany
³⁹ IFAE, Edifici Cn., Campus UAB, E-08193 Bellaterra, Spain
⁴⁰ Yerevan Physics Institute, AM-375036 Yerevan, Armenia
⁴¹ Depto. de Astrofísica, Universidad, E-38206 La Laguna, Tenerife, Spain
⁴² Università di Udine, and INFN Trieste, I-33100 Udine, Italy
⁴³ Institut de Ciències de l'Espai (IEEC-CSIC), E-08193 Bellaterra, Spain
⁴⁴ Inst. de Astrofísica de Andalucía (CSIC), E-18080 Granada, Spain
⁴⁵ University of California, Davis, CA-95616-8677, USA
⁴⁶ Tuorla Observatory, Turku University, FI-21500 Piikkiö, Finland
⁴⁷ Institute for Nuclear Research and Nuclear Energy, BG-1784 Sofia, Bulgaria
⁴⁸ INAF/Osservatorio Astronomico and INFN, I-34143 Trieste, Italy
⁴⁹ ICREA, E-08010 Barcelona, Spain
⁵⁰ Università di Pisa, and INFN Pisa, I-56126 Pisa, Italy
⁵¹ Fred Lawrence Whipple Observatory, Harvard-Smithsonian Center for Astrophysics, Amado, AZ 85645, USA

⁵² Department of Life and Physical Sciences, Galway-Mayo Institute of Technology, Dublin Road, Galway, Ireland

⁵³ Department of Physics and Astronomy and the Bartol Research Institute, University of Delaware, Newark, DE 19716, USA

⁵⁴ Department of Physics and Astronomy, University of California, Los Angeles, CA 90095, USA

⁵⁵ Department of Physics, Washington University, St. Louis, MO 63130, USA

⁵⁶ School of Physics and Astronomy, University of Leeds, Leeds, LS2 9JT, UK

⁵⁷ Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138, USA

⁵⁸ Argonne National Laboratory, 9700 S. Cass Avenue, Argonne, IL 60439, USA

⁵⁹ School of Physics, University College Dublin, Belfield, Dublin 4, Ireland

⁶⁰ School of Physics, National University of Ireland, Galway, Ireland

⁶¹ Astronomy Department, Adler Planetarium and Astronomy Museum, Chicago, IL 60605, USA

⁶² Physics Department, McGill University, Montreal, QC H3A 2T8, Canada

⁶³ Physics Department, University of Utah, Salt Lake City, UT 84112, USA

⁶⁴ Department of Physics, Purdue University, West Lafayette, IN 47907, USA

⁶⁵ Department of Physics, Grinnell College, Grinnell, IA 50112-1690, USA

⁶⁶ Department of Physics and Astronomy, Barnard College, Columbia University, NY 10027, USA

⁶⁷ Santa Cruz Institute for Particle Physics and Department of Physics, University of California, Santa Cruz, CA 95064, USA

⁶⁸ Enrico Fermi Institute, University of Chicago, Chicago, IL 60637, USA

⁶⁹ Department of Physics and Astronomy, Iowa State University, Ames, IA 50011, USA

⁷⁰ Department of Physics and Astronomy, University of Iowa, Van Allen Hall, Iowa City, IA 52242, USA

⁷¹ Department of Physics and Astronomy, DePauw University, Greencastle, IN 46135-0037, USA

⁷² Department of Physics, Pittsburg State University, 1701 South Broadway, Pittsburg, KS 66762, USA

⁷³ Casilla de Correo 67, Sucursal 28, (C1428ZAA) Ciudad Autónoma de Buenos Aires, Argentina

⁷⁴ Kavli Institute for Cosmological Physics, University of Chicago, Chicago, IL 60637, USA

⁷⁵ Department of Applied Physics and Instrumentation, Cork Institute of Technology, Bishopstown, Cork, Ireland

⁷⁶ Laboratoire Leprince-Ringuet, Ecole Polytechnique, CNRS/IN2P3, F-91128, Palaiseau, France

^a deceased

* AGILE Team Corresponding Author: I. Donnarumma, immacolata.donnarumma@iasf-roma.inaf.it

** GASP-WEBT Team Corresponding Author: M. Villata, villata@oato.inaf.it, C. M. Raiteri, raiteri@oato.inaf.it

*** MAGIC Collaboration Corresponding Author: R. M. Wagner, robert.wagner@mpp.mpg.de; C. C. Hsu, cchsu@mpp.mpg.de; K. Satalecka, satalek@ifh.de

**** VERITAS Team Corresponding Author: J. Grube, jeffrey.grube@ucd.ie

Fossati et al. 1998). The first hump is commonly interpreted as due to synchrotron radiation from high-energy electrons in a relativistic jet, while the origin of the second peak is still uncertain. In the leptonic scenario it is interpreted as inverse Compton (IC) scattering of the synchrotron (Synchrotron self-Compton, SSC) or external photons (External Compton, EC) by the same population of relativistic electrons. The observed correlated variability between X-rays and TeV γ -rays (Maraschi et al. 1999; Fossati et al. 2008, Wagner 2008) is well explained in the SSC framework (Ghisellini et al. 1998), whereas the EC scenario is unlikely to apply in HBLs, due to the low density of ambient photons. Alternatively, hadronic models invoke proton-initiated cascades and/or proton-synchrotron emission (Aharonian 2000, Mücke et al. 2003).

Leptonic and hadronic scenarios for HBLs predict different properties of the γ -ray emission in relation to emissions in other energy bands. Specifically, the hadronic models (as opposed to the SSC ones) predict a flatter slope of the ~ 100 MeV IC emission than that of the synchrotron emission in the optical-UV energy bands. γ -ray observations of flaring BL Lacs and simultaneous multiwavelength data are thus the keys to investigating these two scenarios.

A hard X-ray flare of Mrk 421 was detected by SuperAGILE on 2008 June 10 (Costa et al. 2008). This detection was later followed by a detection in γ -rays (Pitorri et al. 2008) by the AGILE/GRID (Gamma Ray Imaging Detector) and prompted a ToO observation by Swift/XRT, complementing the ongoing multifrequency observing campaign of Mrk 421 with WEBT (optical), MAGIC and VERITAS (TeV). We report on the observations and the analysis of these data, complemented by the publicly-available data from RossiXTE/ASM (2-12 keV) and Swift/BAT (15-50 keV), and discuss the spectral energy distribution of the source during this bright γ -ray flare.

2. OBSERVATIONS AND RESULTS

2.1. AGILE observations

AGILE (Tavani et al. 2008) is a small mission of the Italian Space Agency for the exploration of the γ -ray sky, operating in a low Earth orbit since 2007 April 23. The AGILE composite payload allows for simultaneous observations in the energy ranges 30 MeV-30 GeV and 20-60 keV over a very wide field of view by means of GRID and the hard X-ray imager SuperAGILE, respectively. Mrk 421 was observed for five days, between 2008-06-09 17:02 UT and 2008-06-15 02:17 UT.

2.1.1. Hard X-ray observations

On 2008 June 10 SuperAGILE detected enhanced hard X-ray emission from Mrk 421. The measured flux in 20-60 keV was found to be above 30 mCrab (24-hour average), almost an order of magnitude larger than its typical flux in quiescence. The source was at $\sim (0, 20)$ degrees in the orthogonal SuperAGILE reference frame (Feroci et al. 2007), thus exposing an effective area $\sim 55\%$ of the full on-axis value. In the following days, the flux increased up to about 55 mCrab. The 5-day 20-60 keV SuperAGILE light curve is shown in Fig. 1c. Using the publicly available light curves for this source

from the BAT experiment¹ in the 15-50 keV energy range (near the SuperAGILE bandpass), we calculated daily averages of the sparse observations. This allowed us to obtain good coverage also before and after the AGILE observation (black squares in Fig. 1c), revealing that SuperAGILE indeed observed the maximum brightness of this hard X-ray flare. Both SuperAGILE and BAT count rates were converted to physical units by assuming a Crab-like spectrum.

SuperAGILE photon-by-photon data allows extraction of a time-averaged energy spectrum from the mask-convolved data. Given the lack of substantial spectral variability in the SuperAGILE/ASM hardness ratio (Fig.1e) we accumulated the average energy spectrum from the data of the last 4 days of the observations, when the source flux varied between 35 and 55 mCrab, for a total net source exposure of ~ 140 ks. Despite poor statistics, this 4-point energy spectrum is able to reasonably constrain the photon index of a simple power law, $\Gamma = 2.43_{-0.64}^{+0.69}$ ($\chi^2_{\text{dof}} = 0.8, 2$ dof). The average flux is $F(20 - 60\text{keV}) = (4.90 \pm 0.54) \times 10^{-10}$ erg cm⁻²s⁻¹ ($(9.8 \pm 1.1) \times 10^{-3}$ photons cm⁻² s⁻¹).

2.1.2. Gamma-ray Observations

The GRID data were analyzed using the AGILE standard pipeline (Vercellone et al. 2008), with a bin-size of $0.25^\circ \times 0.25^\circ$. Only events flagged as γ -rays and not recorded while the satellite crosses the South Atlantic Anomaly were accepted. We also rejected all events with reconstructed direction within 10° from the Earth limb thus reducing contamination from Earth's γ -ray albedo.

Mrk 421, observed $\sim 20^\circ$ off-axis with respect to the boresight, was not detected on daily time scales, implying a daily-averaged flux smaller than $\sim 100 \times 10^{-8}$ photons cm⁻² s⁻¹, similar to what observed by EGRET (Hartman et al. 1999). A 4.5- σ significance in the range 100 MeV-10 GeV resulted from an integration over the whole 5-day period (net exposure ~ 260 ks). The measured flux is $(42_{-12}^{+14}) \times 10^{-8}$ photons cm⁻² s⁻¹, about ~ 3 times higher than the average flux detected by EGRET ($\sim 13 \times 10^{-8}$, (Hartman et al. 1999)) and ~ 1.5 times higher than, but still consistent, with the highest flux $(27 \pm 7) \times 10^{-8}$ photons cm⁻² s⁻¹ observed during the Viewing Period 326. The short duration of the AGILE observation did not allow us to follow the possible decay of the γ -ray emission. Due to the relative faintness of the source and the short exposure, the GRID data do not allow us to extract a reliable energy spectrum.

2.2. The Soft X-ray band

2.2.1. Swift X-ray telescope

Following the SuperAGILE detection, on 2008 June 12 we triggered a ToO observation with the Swift X-Ray Telescope (XRT) that promptly observed the source for 5ks between 2008 June 12 19:33:20 UT (MJD 54629.8) and June 13 at 01:57:37 UT (MJD 54630.1).

The XRT data were processed using the standard procedure (`xrtpipeline` v0.12.0) developed at the ASI Science Data Center and distributed by the NASA/HEASARC. Observations were taken in Window Timing mode at a count rate of 120–150 cts s⁻¹. We

¹ <http://swift.gsfc.nasa.gov/docs/swift/results/transients/weak/Mrk421/>

selected XRT grades 0–2 (Burrows et al. 2005) and extracted the WT data in a rectangular 40×20 -pixel region centered on the source. The background was also extracted within a box (40×20 -pixel), far from background sources.

Due to calibration uncertainties, we restricted our spectral fit to the energy range 0.7–9.0 keV and added a 3% systematics to the model (Cusumano et al. 2006). The XRT spectral data are well described by an absorbed log-parabolic model. A joint fit of the XRT and SuperAGILE spectral data using the 4-day average spectrum described in Sect. 2.1.1. used the log-parabolic model with Galactic absorption ($N_{\text{H}}^{\text{Gal}} = 1.61 \times 10^{20} \text{ cm}^{-2}$; Lockman & Savage 1995), defined as:

$$F(E) = K E^{-a-b \log(E)} \text{ photons cm}^{-2} \text{ s}^{-1}$$

where a is the photon index at 1 keV and b is the curvature parameter (Massaro et al. 2004; Massaro et al. 2008). This model usually describes adequately the featureless and curved spectrum in HBLs. The joint fit provides $a = 1.65^{+0.01}_{-0.02}$, $b = 0.37^{+0.01}_{-0.005}$ ($\chi^2_{\text{dof}}=1.4$, 763 dof), which implies a peak energy $2.97^{+0.22}_{-0.18}$ keV, and predicts $F_{2-10\text{keV}} = 2.56 \times 10^{-9} \text{ erg cm}^{-2} \text{ s}^{-1}$ (or $0.4 \text{ photons cm}^{-2} \text{ s}^{-1}$) and $F_{20-60\text{keV}} = (5.7 \pm 0.6) \times 10^{-10} \text{ erg cm}^{-2} \text{ s}^{-1}$ ($(1.1 \pm 0.1) \times 10^{-2} \text{ photons cm}^{-2} \text{ s}^{-1}$), comparable to the stand-alone SuperAGILE best fit.

2.2.2. RossiXTE All Sky Monitor (ASM)

Given the high flux observed by Swift/XRT, we retrieved the public light curves provided by the ASM² to trace the evolution of the soft X-rays during the AGILE observation. Fig. 1b shows the daily light curve of Mrk 421 in the energy range 2–12 keV, obtained by properly averaging the dwell-by-dwell data.

The emission at soft X-rays is well correlated with hard X-ray emission. The ASM data show that the XRT observation took place when the source was at its maximum emission at soft X-rays (MJD ~ 54630). Comparing the relative intensity of the two flares in Fig. 1, it appears that the second peak is significantly harder than the first one. This is also shown in Fig. 1e, where we computed the daily-averaged hardness ratio between hard (15–60 keV) and soft (2–12 keV) X-rays. The source appears to have undergone the hardest part of this double-humped flare just during the AGILE γ -ray detection.

2.3. Observations in the VHE γ -rays

The 4-telescope array VERITAS (Acciari et al. 2008) at the Fred Lawrence Whipple Observatory (Arizona) and the single-dish instrument MAGIC (Baixeras et al. 2004; Tesaro et al. 2007) at La Palma are imaging air Cherenkov telescopes covering an energy range from ~ 100 GeV to some tens of TeV. The instruments have a typical energy resolution of $<20\%$ (VERITAS) and $20\text{--}30\%$ (MAGIC), and event-by-event angular resolution of $<0.14^\circ$. Wobble-mode observations (Daum et al. 1997) at an 0.4° offset from the camera center were taken on five nights from 2008 June 3–8 with MAGIC at zenith angles (ZA) between 28° and 48° and on 4 nights (May 27, June 1, 5, 6) with VERITAS (wobble offset: 0.5°) at




FIG. 1.— **a)** R -band optical light curve from GASP-WEBT (May 24–June 23); **b)** ASM (2–12 keV) light curve (bin size is 1 day) and XRT (2–10 keV) flux (blue triangle); **c)** SuperAGILE (20–60 keV, blue triangles; 1 Crab = $0.2 \text{ ph cm}^{-2} \text{ s}^{-1}$) and BAT (15–50 keV, empty black squares; 1 Crab = $0.29 \text{ ph cm}^{-2} \text{ s}^{-1}$); **d)** MAGIC and VERITAS (>400 GeV, empty black squares and black circles, respectively), the Crab flux >400 GeV (horizontal dashed line), AGILE (>100 MeV, blue triangle); **e)** the hardness ratio computed by using the SuperAGILE and ASM data for each day. The dashed vertical lines mark period 1 and period 2 (see text for details).

ZA between 32° and 40° during partial moon light conditions. The total live-time after applying quality selection is 2.95 and 1.17 hours with MAGIC and VERITAS, respectively. The data were analyzed using the MAGIC (Albert et al. 2008a) and VERITAS (Daniel et al. 2007; Acciari et al. 2008) standard calibration and analyses and image parameters (Hillas 1985). For VERITAS, the γ -ray direction and air shower impact parameter on the ground are reconstructed using the stereoscopic techniques in Hofmann et al. (1999) and Krawczynski et al. (2006). Any γ -ray excess is derived from the θ^2 distribution, where θ represents the angular distance between

² <http://xte.mit.edu/asmlc/ASM.html>

the source position in the sky and the reconstructed arrival position of the air shower. For MAGIC, θ is estimated using the DISP method (Fomin et al. 1994). For VERITAS, 99.9% of the background of cosmic-rays is rejected by using selection cuts on θ^2 , mean scaled width and length, and by using quality cuts of each event. The MAGIC analysis utilizes a random forest method (Albert et al. 2008b) to discriminate the dominating background of hadronic cosmic-ray events and for the energy estimation of the γ -ray events. The energy and effective area of each event is reconstructed from Monte Carlo simulations. The integral flux and energy spectrum of the source is then derived from the effective areas for nights with a significant detection. Two independent analyses of both the MAGIC and VERITAS data sets yielded consistent results. In total, a signal corresponding to a significance level of 44σ (VERITAS) and 66σ (MAGIC) is obtained by following eq. 17 of Li & Ma (1983). The combined MAGIC-VERITAS data (Fig. 1d) show a transient peaking near MJD 54622. The VERITAS energy spectrum for June 6 is provided. A power law fit over the energy range 0.3–5 TeV resulted in a $\chi^2_{\text{dof}}=0.7$ with a photon index $\Gamma = 2.78 \pm 0.09$. In Fig. 2 we show the intrinsic γ -ray spectrum at the source reconstructed by removing attenuation effects by the extragalactic background light (Hauser & Dwek 2001) following the procedure of Raue & Mazin (2008) ⁽³⁾. Fitting a power law to the intrinsic spectrum yields a photon index $\Gamma = 2.59 \pm 0.08$, which is not significantly harder than the measured spectrum due to the relatively low redshift $z = 0.031$.

2.4. Optical and UV observations

Mrk 421 is one of the 28 γ -ray-loud blazars that are regularly monitored by the GLAST-AGILE Support Program (GASP; Villata et al. 2008) of the Whole Earth Blazar Telescope (WEBT)⁴. The GASP observations started in early 2007 November. The *R*-band data were calibrated according to the photometric sequence by Villata et al. (1998). A careful data analysis is warranted because the source flux is contaminated by the emission of the host galaxy and nearby objects. The flux contribution of these objects was subtracted according to Nilsson et al. (2007). Moreover, we corrected for a Galactic extinction of $A_R = 0.042$ mag. The resulting *R*-band light curve⁵ during the period considered in this paper is shown in Fig. 1a.

During the Swift pointing on 2008 June 12–13, the UVOT (Poole et al. 2008) instrument observed Mrk 421 in the UVW1 and UVW2 photometric bands. The UVOTSOURCE tool is used to extract counts, correct for coincidence losses, apply background subtraction, and calculate the source flux. We applied a standard 5 arcsec radius source aperture, and a 20 arcsec background region. The source fluxes are dereddened using the interstellar extinction curve in Fitzpatrick (1999).

³ <http://www.desy.de/~mraue/eb1/>

⁴ <http://www.oato.inaf.it/blazars/webt>

see e.g. (Villata et al. 2007; Raiteri et al. 2007)

⁵ The optical data presented in this paper are stored in the GASP-WEBT archive; for questions regarding their availability, please contact the WEBT President Massimo Villata villata@oato.inaf.it.

3. DISCUSSION

Mrk 421 showed a very interesting broad-band activity during the first half of 2008 June as derived from AGILE data combined with those of WEBT, Swift, MAGIC and VERITAS. Using our multi-frequency data we were able to derive time-resolved SEDs (Fig. 2). We distinguish two time periods: *period 1*: 2008 June 6, with the inclusion of optical, X-ray (XTE and BAT) and TeV data (VERITAS); and *period 2*: 2008 June 9–15, including optical, UV, X-rays (XRT and SuperAGILE) and gamma-ray data (AGILE). The source shows a very interesting time-variable broad band emission that appears to be in overall agreement with an SSC model. The optical, soft and hard X-ray bands strongly constrain the SED around the synchrotron peak, and its daily variability reveals the physical processes of Mrk 421. Possible correlated variability is shown in Fig. 1 between the optical (an overall decreasing trend with superimposed spikes of emission), the X-rays (several emission peaks lasting few days), and the high-energy parts of the spectrum. Based on the physical constraints obtained for the synchrotron peak, we can model both the HE and VHE γ -ray emission. The data collection and broad-band SED extends over 12 decades in energy. Taking advantage of the overlapping MAGIC and VERITAS observations, we present a combined VHE light curve using the current generation of Northern imaging air Cherenkov systems.

We first model the synchrotron peak of emission using the period 1 optical, soft and hard X-ray data. The short time-variability (Fig. 1) constrains the size of the emitting region to $R < cT\delta \sim 5 \times 10^{16} (\delta/20)$ cm. Hence, we consider a one-zone SSC model (Tavecchio et al. 1998) based on a blob of comoving size $R = 4 \times 10^{16}$ cm, with a relativistic Doppler factor $\delta = 20$ and characterized by non-thermal comoving electron energy distribution function described by a double power law:

$$n_e(\gamma) = \frac{K\gamma_b^{-1}}{(\gamma/\gamma_b)^{p_1} + (\gamma/\gamma_b)^{p_2}} \quad (1)$$

where the comoving Lorentz factor (γ) varies in the range $\gamma_{\text{min}} = 4 \times 10^3 < \gamma < \gamma_{\text{max}} = 1.3 \times 10^6$, the normalization (density) constant $K = 4 \times 10^{-4} \text{ cm}^{-3}$, and the break energy $\gamma_b = 3.6 \times 10^5$ and with $p_1 = 2.22$, $p_2 = 4.5$, the low-energy and high-energy power-law indices, respectively (see Table 1). With these parameters we found that the data for period 1 are best fitted with a comoving magnetic field $B = 0.1$ G.

Variability may be caused by several factors; we consider two cases: (A) hardening/softening of the electron energy distribution function caused by particle acceleration processes; (B) increase/decrease of the comoving particle density, as a consequence of additional particle injection/loss by shock processes.

We expect TeV variability to be comparable with the X-ray one if case (A) applies: this is because the emission is in the Klein-Nishina regime. Alternately, for case (B) we expect the TeV relative variability ($\Delta F/F$) to be a factor of 2 greater than that of the X-ray flux variability.

Our AGILE, MAGIC and VERITAS data appear to support case (A). We compare the SEDs for period 1 and period 2, to better assess the spectral evolution. In Fig. 2 we show our optimized modelling of the time-resolved synchrotron peak and consequent SSC high-energy emission for the period 1 as well as for 2008 June 12–13 of the

period 2. In the last case, the adopted model parameters are $p_1 = 2.1$, $p_2 = 5$, $\gamma_b = 4.2 \times 10^5$, $K = 6 \times 10^{-4} \text{ cm}^{-3}$. Our model predicts an even larger TeV flux for period 2 (no TeV observations exist, however) than detected in period 1.

A detailed discussion of the complex optical vs. X-ray variability of Mrk 421 as shown in Fig. 1 will be presented elsewhere. We notice here a few remarkable points. The optical light curve shows variations of the order of 10% on a time scale \sim few days, superimposed on a long decay during the entire period. Individual soft and hard X-ray peaks result in increased fluxes by a factor of ~ 2.5 and ~ 5 , respectively: no long term decay appears. This different behaviour of the X-ray radiation and the bulk of the optical emission may interestingly suggest a more complex scenarios than A) and B) ones: optical and X radiation comes from two different jet regions, each one characterized by its own variability. A possible scenarios is one in which the inner jet region would produce the X-rays and it would be at least partially transparent to the optical radiation. In contrast, the outer region can only produce lower-frequency emission. The signature of the X-ray events visible in the optical light curve would come from the inner region and would be diluted by the optical radiation emitted from the outer region (see Villata & Raiteri 1999 for the case of Mrk 501; Villata et al. 2004).

Interestingly, the 2-10 keV flux measured by XRT on June 12 – 13, $\sim 2.6 \times 10^{-9} \text{ erg cm}^{-2} \text{ s}^{-1}$, is higher than all previous observations ($< 2 \times 10^{-9} \text{ erg cm}^{-2} \text{ s}^{-1}$; Fosfati et al. 2008; Lichti et al. 2008). A joint analysis of the XRT and SuperAGILE data, covering the range from 0.7 to 60 keV, provides a best-fit spectral model

consistent with a log-parabolic shape, with parameters implying a peak energy $\sim 3 \text{ keV}$, in good agreement with the steeper positive correlation between the peak energy and the maximum of the SED found by Tramacere et al. 2007 (see their Fig. 3), although our value of the peak energy shows a significant shift with respect to typical values of 0.5-1 keV for this source.

We conclude our analysis of the broad-band variable emission from Mrk 421 by emphasizing that our multi-telescope/instrument data show a very interesting variability that provides support for an SSC model of the source. The γ -ray emission detected by AGILE during period 2 and the TeV emission detected during period 1 can be successfully modelled from the characteristics of the corresponding synchrotron peaks.

AGILE is a mission of ASI, with co-participation of INAF and INFN. This work was partially supported by ASI grants I/R/045/04, I/089/06/0, I/011/07/0 and by the Italian Ministry of University and Research (PRIN 2005025417), (ASDC) I/024/05/1. The MAGIC collaboration thanks the Instituto de Astrofísica de Canarias for the excellent working conditions at the Observatorio del Roque de Los Muchachos in La Palma and support by the German BMBF and MPG, the Italian INFN and Spanish MCINN is acknowledged. This work was supported by ETH Research Grant TH 34/043, by the Polish MNiSzW Grant N N203 390834, the YIP of the Helmholtz Gemeinschaft, the grant of Georgian National Science Foundation GNSF/ST07/4-180. E. K. acknowledges financial support from the NCS grant No. 96-2811-M-008-058. The VERITAS collaboration is supported by grants from the U.S. Department of Energy, the National Science Foundation, and the Smithsonian Institution, by NSERC in Canada, Science Foundation Ireland, and PPARC in the UK. We acknowledge the technical support staff at the FLWO. We also acknowledge the Swift Team for carrying out the ToO observation.

REFERENCES

- Acciari, V. A. et al. (VERITAS Collab.) 2008, *ApJ*, 679, 1427
 Aharonian, F. A. 2000, *NewA*, 5, 377
 Albert et al. (MAGIC Collab.) 2008a, *ApJ*, 674, 1037
 Albert et al. (MAGIC Collab.) 2008b, *Nucl. Instr. Meth. A* 588, 424
 Baixeras, C. et al., 2004, *Nucl. Instrum. Meth. A* 518, 188
 Burrows, D. N., Hill, J. E., Nousek, J. A., et al. 2005, *Space Sci. Rev.*, 120, 165
 Costa, E., Del Monte, E., Donnarumma, I., et al. 2008, ATEL #1574
 Cusumano, G. in: *Proc. SWIFT and GRBs: Unveiling the Relativistic Universe*, June 5-9, 2006, Venice (Italy)
 Daniel, M. K. et al. 2007, *Proc. 30th ICRC*, Merida, Mexico, 283
 Daum A. et al. (HEGRA Collab.) 1997, *Astropart. Phys.*, 8, 1
 Feroci, M., Costa, E., Soffitta, P., et al. 2007, *Nucl. Instrum. Methods Phys. Res. A*, 581, 728
 Fitzpatrick, E., 1999, *PASP*, 111, 63
 Fomin V.P. et al. 1994, *APh*, 2, 137
 Fosfati, G., Maraschi, L., Celotti, A. et al. 1998, *MNRAS*, 299, 433
 Fosfati, G., Buckley, J. H., Bond, I. H., et al. 2008, *ApJ*, 677, 906
 Ghisellini, G., Celotti, A., Fosfati, G., et al. 1998, *MNRAS*, 301, 451
 Hartman, R. C., Bertsch, D. L., Bloom, S. D., et al. 1999, *ApJS*, 123, 79
 Hauser, M. G., Dwek, E., 2001, *ARA&A*, 39, 249
 Hillas A.M. 1985, in *Proc. 19th ICRC*, La Jolla, USA, 3, 445
 Hofmann, W. et al. 1999, *Astropart. Phys.*, 12, 135
 Holder, J. et al. (VERITAS Collab.) 2006, *Astropart. Phys.*, 25, 361
 Krawczynski, H. et al. 2006, *Astropart. Phys.*, 25, 380
 Lichti, G. G., Bottacini, E., Ajello, M., et al. 2008, *A&A*, 486, 721
 Lin, Y. C., Bertsch, D. L., Chiang, J., et al. 1992, *ApJ*, 401, 61
 Lockmann, F. J., & Savage, B. D. 1995, *ApJS*, 97, 1
 T.-P. Li and Y.-Q. Ma, 1983, *ApJ*, 272, 317.
 Maraschi, L., Fosfati, G., Tavecchio, F. et al. 1999, *ApJ*, 526L, 81M
 Massaro, E., Perri, M., Giommi, P. et al. 2004, *A&A*, 422, 103




Fig. 2.— SEDs of Mrk 421 obtained by combining the GASP-WEBC, SWIFT/UVOT, RossiXTE/ASM, XRT, SuperAGILE, BAT, GRID and VERITAS data in period 1 and period 2 (red empty circles and black filled circles, respectively). Both are one-zone SSC models (red dashed line for period 1 and black solid line for period 2).

- Massaro, F., Tramacere, A., Cavaliere, A., Perri, M. & Giommi, P. 2008, *A&A*, 478, 395
 Mücke, A., Protheroe, R. J., Engel, R., et al. 2003, *APh*, 18, 593

TABLE 1
SSC MODEL PARAMETERS.

parameter	period 1	period 2
γ_b	$3.6 \cdot 10^5$	$4.2 \cdot 10^5$
γ_{max}	$1.3 \cdot 10^6$	$1.3 \cdot 10^6$
p_1	2.22	2.1
p_2	4.5	5
B [G]	0.1	0.1
K [cm ⁻³]	$4 \cdot 10^{-4}$	$6 \cdot 10^{-4}$
δ	20	20
θ [°]	2	2

Nilsson, K., Pasanen, M., Takalo, L. O., et al. 2007, A&A, **475**, 199
 Padovani, P. & Giommi, P. 1995, ApJ, 444, 567
 Petry, D., Ishibashi, W., Beckmann, V., et al. 2008, ATEL #1507
 Pittori, C., Cutini, S., Gasparrini, D., et al. 2008, ATEL #1583
 Poole, T. S. et al. 2008, MNRAS, 383, 627
 Punch, M., Akerlof, C. W., Cawley, M. F., et al. 1992, Nature, **358**, 477
 Raiteri C. M., Villata, M., Larionov, V. M., et al. 2007, A&A, **473**, 819

Raue M., Mazin, D. 2008, Int. J. Mod. Phys. D, 17, 1515
 Sambruna, R. M. and Maraschi, L. and Urry, C. , 1996, ApJ, **463**, 444
 Tavani et al. 2008, arXiv:0807.4254v1
 Tavecchio, F., Maraschi, L., Ghisellini, G. et al. 1998, ApJ, 509, 608
 Tescaro, D., Bartko, H., Galante, N., et al. (MAGIC Collaboration) 2007, in: Proc. 30th ICRC, Merida, Mexico (preprint: arXiv:0709.1410)
 Tramacere, A., Giommi, P., Massaro, E. et al., 2007, A&A, 467, 501
 Vercellone, S., Chen, A. W., Giuliani, A., et al. 2008, ApJ, **676**, 13
 Villata, M., Raiteri, C. M., Lanteri, L., et al. 1998, A&AS, **130**, 305
 Villata, M., & Raiteri, C. M. 1999, A&A, 347, 30
 Villata, M., Raiteri, C. M., Kurtanidze, O. M., et al. 2004, A&A **421**, 103
 Villata, M., Raiteri, C. M., Aller, M. F., et al. 2007, A&A, **464**, L5
 Villata, M., Raiteri C. M., Larionov, V. M et al. 2008, A&A, **481**, L79
 Wagner, R.M., 2008, PoS(BLAZARS2008), 63, 013 (arXiv:0809.2843)